

**POSTER PRESENTATION****Open Access**

# Diffuse tensor cardiac MRI evaluation of fiber architecture of athlete hypertrophic heart in vivo

Ming-Ting Wu<sup>1\*</sup>, Wen-Yih I Tseng<sup>2,3</sup>, Mao-Yuan M Su<sup>3</sup>, Van J Weeden<sup>4</sup>, Timothy G Reese<sup>4</sup>From 15th Annual SCMR Scientific Sessions  
Orlando, FL, USA. 2-5 February 2012**Background**

The fiber architecture adaption in physiological hypertrophy of the athlete heart is still elusive. We aimed to use diffusion tensor cardiac MR (DT-CMR) to evaluate the tissue property and fiber architecture of elite athlete heart.

**Methods**

Eight elite athletes of Marathon runner (endurance-training type), 8 of weight-lifter (strength-training type) and 8 ordinary style (medical interns) were enrolled. Each subject received a CMR study on a 1.5 T scanner including 1. cine SSFP of a stack of LV short axis for LV mass and function; 2 DT-CMR, ECG-gated stimulated echo diffuse EPI on three levels of LV. Diffuse tensor composed of 6 directions and b value = 300 mm<sup>2</sup>/sec. 3. phase-contrast flow measurement at ascending aorta for stroke volume. The data were compared between groups and correlated between the parameters.

**Results**

The myocardium showed no difference of mean diffusivity (MD) and fractional anisotropy between the groups. Weight lifter showed increase of stroke volume / BSA and LV mass / BSA as compared to runner and ordinary groups. The fiber architecture showed an increased proportion of right-handed helical fibers (mainly in the sub-endocardial zone) in runner and lifter equally, as compared to ordinary group. Putting all 24 subjects together, there was a linear regression between the proportion of right-handed helical fiber and LV mass ( $R^2 = 0.38$ ,  $p = 0.002$ ).

**Conclusions**

DT-CMR revealed the physiological hypertrophy of athlete heart was mainly due to right-handed helical fibers. This underscores the important role of subendocardial fiber on the LV function.

**Funding**

N/A

**Author details**<sup>1</sup>Department of Radiology, National Yang Ming University, Taipei, Taiwan.<sup>2</sup>Center for Optoelectronic Biomedicine, National Taiwan University, Taipei, Taiwan. <sup>3</sup>Department of Medical Imaging, National Taiwan University Hospital, Taipei, Taiwan. <sup>4</sup>National Yang Ming University, Taipei, Taiwan.

Published: 1 February 2012

doi:10.1186/1532-429X-14-S1-P170


**Cite this article as:** Wu et al.: Diffuse tensor cardiac MRI evaluation of fiber architecture of athlete hypertrophic heart in vivo. *Journal of Cardiovascular Magnetic Resonance* 2012 **14**(Suppl 1):P170.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

<sup>1</sup>Department of Radiology, National Yang Ming University, Taipei, Taiwan  
Full list of author information is available at the end of the article



**Figure 1** Regression between LV mass and right-handed helical fiber percentage across the 24 subjects.